

Evaluation of some commercial formulations against *Spodoptera littoralis* and *Hypera brunneipennis* larvae.

Narmen A. Youssef and Atef A. Abd-ELgayed

Dept. of Plant Protection, Fac. of Agriculture, Fayoum University, Fayoum, Egypt.

Abstract

The insecticidal activity of three commercial Bt – formulations and one fungus; namely DipelDf , W- Bus and Protecto (*Bacillus thuringiensis var.kurstaki*) and Biofly (*Beauveria bassiana*) were tested against 2nd and 4th instar larvae of *Spodoptera littoralis* (Boisd) and *Hypera brunneipennis* (Boheman) were studied under laboratory conditions. Results revealed that Bt- formulations caused the larval mortality after treatment of *S. littoralis* 2nd and 4th larval instars ranged from 40 to 100 % and 32.5 to 92.5 % and reached 100 ,85 and 100 % for *H. brunneipennis*, respectively at the highest concentration after 7 days of treatment compared to 77.5 % and 60 % for *S. littoralis* and 95 and 85 % for *H. brunneipennis* treatments by Biofly. The DipelDf and W-Bus were highly efficient on the insect larvae, followed by Biofly and Protecto, respectively. Based on the LC₅₀ values, DipelDf was the highest toxic to *S. littoralis* and *H. brunneipennis* than that of the other compounds. Pupation and adults emergence percentages were reduced by all treatments compared to control.

Key words: *Spodoptera littoralis* ,*Hypera brunneipennis*, commercial products , efficiency

Introduction

Among chewing insect pests , *Spodoptera littoralis* (Boisd) (Lepidoptera: Noctuidae) is considered as an important sporadic pest in the world. It causes 25-100 % economic loss (Dhir *et al.*, 1992; Prayogo *et al.*, 2005) in crops based on crop stage and its population level in the field. The Egyptian alfalfa weevil (EAW) , *Hypera brunneipennis* (Boheman) (Coleoptera : Curculionidae) is considered to be the most serious and destructive pest of alfalfa in Egypt (Al-Doghairi and Elhag, 2003).One annual generation is recorded in Egypt for the EAW (Hammad *et al.*, 1967). The larval stage is the most damaging during the weevil life cycle. By feeding on the alfalfa plants growing tips, the larvae cause skeletonization of leaves, stunting, reduced plant growth, and ultimate reduction in yield. The adults are also, foliar feeders, causing additional, but less significant, damage.

The widespread and intensive use of different synthetic insecticides for controlling this pest increased environmental problems such as insect resistance, excessive persistence of residues, human health hazards and harmful effect on the non- target organisms. From this point of view, it is necessary to minimize the application of pesticides that considered as a main source of environmental pollution and use other compounds may proof as good alternatives of insecticides. In recent years, crop protection based on biological control of crop pests with microbial pathogens as virus, bacteria, fungi and nematodes were considered as valuable tools in pest management (Bhattacharya *et al.*, 2003). Entomopathogenic fungi may proof, also, as valuable and play an important role in integrated pest management programs. (El- Hawary and Abd El-

Salam, 2009) reported that fungal biological control agents have demonstrated efficacy against a wide range of insect pests including *S. litura* . Successful use of fungal pathogens in pest control depends on selection of right virulent fungal strain formulated in proper way and applied at an appropriate dose against susceptible host stage under favorable environmental conditions (Asi *et al.*, 2012).Among the entomopathogenic agents, also, the most widely used biopesticides are subspecies and strains of *Bacillus thuringiensis* (Bt). *B. thuringiensis* is a spore-forming bacterium well- known for its insecticidal properties due to its ability to produce crystal inclusions during sporulation. Each strain of this bacterium specifically kills one or a few related species of insect larvae such as Lepidopteran, Dipteran and Coleopteran (Haggag, 2013). Commercial *Bt* products, generally, consist of a mixture of spores and crystals, produced in large fermenters and applied as foliar sprays, much like synthetic insecticides (Sanchis *et al.*, 1999). It is known that most Bt formulations have a very short residual activity.The persistence of Bt. spores show an obvious reduction after few days of exposure to weather, and reduction in its viability was progressively correlated with the time elapsed after exposure in the field. The pathogen is not mobile and cannot escape under the unfavorable conditions. (Mohamed *et al.*, 2010) .

In the present experiments, the effectiveness of several bioinsecticides against the cotton leafworm *S. littoralis* and alfalfa weevil, *H. brunneipennis* was determined with the intention to find out the best compounds for controlling these economic pests in an integrated pest management program.

Material and Methods

Tested insects:**A- *S.littoralis***

The cotton leafworm larvae of *S.littoralis* were obtained from Agricultural Research Centre, Cairo, Egypt, and were reared on fresh leaves of castor bean (*Ricinus communis*) under laboratory conditions of $25 \pm 2^\circ\text{C}$ and $65 \pm 5\%$ R.H.. (Adham *et al.*, 2009 and Kamel *et al.*, 2010). As larvae reached the 2nd and 4th instars, they were used in the experiments.

B- *H.brunneipennis*

Alfalfa weevil larvae were collected, early in the morning, by using an insect sweepnet in an alfalfa field at Fayoum Government. Insects, were reared on fresh alfalfa plants (*Medica gossypifolia* L.) at laboratory conditions of $25 \pm 2^\circ\text{C}$, $65 \pm 5\%$ R.H. and 2nd and 4th instars of the weevils larvae were selected for experiments.

Tested compounds

Commercial formulations of the following insecticides tested against 2nd and 4th instar larvae of *S.littoralis* and *H. brunneipennis* were obtained from the Agricultural Research Centre, Cairo, Egypt. DipelDf (WP) 6.4% : commercial product formulation contains 32×10^3 IU/mg of *Bacillus thuringiensis* var. *kurstaki*; W-Bus (WP) 8% : commercial product formulation contains 8×10^3 IU/mg of *Bacillus thuringiensis* var. *kurstaki*, Protecto (WP) 9.4%: commercial product formulation contains 32×10^6 IU/mg of *Bacillus thuringiensis* var. *kurstaki* and Biofly (WP) : commercial product formulation contains 30×10^6 spores/mg of *Beauveria bassiana*.

Bioassay

The insecticidal activities of the tested Bt-formulations and fungi, each at four concentrations were prepared in distilled water and tested against 2nd and 4th instar larvae of *S.littoralis* and *H. brunneipennis* larvae using the dipping leaf technique (Ahmed, 2009). The leaves were first washed with distilled water and dipped in solution of the desired concentration of Bt or fungi commercial formulations (DipelDf, W-Bus, Protecto and Biofly). Each leaf was dipped for 30 seconds, then placed individually in Petri-dishes (9 cm diameter) containing moistened filter papers to avoid desiccation of leaves. Other castor bean leaves for treatment of *S. littoralis* and alfalfa for *H. brunneipennis* were treated with sterile distilled water for control. Then, ten larvae from each 2nd/ or 4th instars larvae were separately placed in each Petri dish for each treatment. Four Petri-dishes were used as replicates for each treatment and control. Larvae

were allowed to feed for 48h. on treated leaves. Then these leaves were removed and replaced by another untreated ones. All Petri-dishes were kept at the above mentioned conditions. Larvae were examined daily for 7 days after treatment to determine the mortality percentages. Accumulative larval mortality was recorded and corrected using Abbott's formula (1925). Afterwards, the corresponding concentration probit lines were estimated in addition to determining 50% mortalities and slope values of tested compounds were also estimated. Data were analyzed by ANOVA and the means were separated using the Duncan's multiple range test (Duncan, 1955).

Fourty newly hatched 2nd, 4th instar larvae of each *S.littoralis* and *H. brunneipennis* were fed as previously described (ten larvae/ four replicates) on leaves treated with the calculated LC_{50} starting of exposure was 2 days after application for each of these compounds. The initial (2 days after application) and residual effect of Bt and fungi formulations at (4, 6 and 8 days) after application against larvae were recorded at the end of the experiment (6) days.

The surviving larvae were transferred to other clean Petri-dishes, and supplied with untreated fresh castor bean leaves until pupation. Pupation and adult emergence percentages after treatment by the LC_{50} , and control were also determined.

Results and Discussion**Toxic effect of Bt and fungal formulations against 2nd and 4th instar larvae of *S.littoralis* and *H. brunneipennis***

Efficacies of the four concentrations of all tested insecticides on 2nd and 4th instar larvae of *S.littoralis* and *H. brunneipennis* at 7 day after treatment are presented in table 1. DipelDf, W-Bus and Protecto caused 100% mortality after treatment by highest concentration on the 2nd instar larvae of *H. brunneipennis* while treatment of 4th instar larvae caused 100, 90 and 85%, respectively and 95, 85% mortality at Biofly. While the larval mortality was in the range 40 to 100 and 32.5 to 92.5% on 2nd and 4th instar larvae of *S. littoralis*, respectively at Bt formulations and 77.5, 60% at Biofly. There were significant differences between the tested insecticides of both insects ($F=75.08$; $df=3$ for *S. littoralis*) and ($F= 21.31$; $df= 3$ for *H. brunneipennis*), respectively at 7 days post treatment. Also, there were significant differences between concentrations and also significant between 2nd and 4th instar larvae of both insect species ($F=38.18$; ; $df=3$ for *S.littoralis*) and ($F= 30.2$; ; $df=3$ for *H. brunneipennis*), respectively.

Table 1. Accumulated corrected mortality percentages after 7 days of treatment by some commercial Bt and fungi formulations against *S. littoralis* and *H. brunneipennis* 2nd and 4th instar larvae.

Formulations	Conc.gm/L	Mortality %			
		<i>Spodoptera littoralis</i>		<i>Hypera brunneipennis</i>	
		2 nd	4 th	2 nd	4 th
Dipel DF	0.5	55 ^{def}	45 ^{cde}	90 ^{ab}	75 ^{cd}
	1	62.5 ^{bcde}	60 ^{bc}	92.5 ^a	82.5 ^{bc}
	2	87.5 ^{ab}	85 ^a	100 ^a	100 ^a
	4	100 ^a	92.5 ^a	100 ^a	100 ^a
	Mean	76.25 ^a	70.63 ^a	95.63 ^a	89.38 ^a
W-Bus	0.5	52.5 ^{defg}	40 ^{cde}	70 ^{bc}	60 ^{efg}
	1	57.5 ^{cdef}	50 ^{cd}	90 ^{ab}	75 ^{cd}
	2	60 ^{cdef}	55 ^{bc}	100 ^a	82.5 ^{bc}
	4	82.5 ^{abc}	77.5 ^{ab}	100 ^a	90 ^{ab}
	Mean	63.13 ^b	55.63 ^b	90 ^{ab}	76.88 ^b
Protecto	0.5	15 ^g	12.5 ^g	60 ^c	50 ^{gh}
	1	22.5 ^{fg}	17.5 ^{fg}	65 ^c	52.5 ^{fgh}
	2	27.5 ^g	25 ^{efg}	70 ^{bc}	65 ^{def}
	4	40 ^{efg}	32.5 ^{defg}	100 ^a	85 ^{bc}
	Mean	26.25 ^c	21.88 ^d	73.75 ^c	63.13 ^c
Biofly	1	40 ^{efg}	30 ^{defg}	70 ^{bc}	45 ^h
	2	50 ^{efg}	35 ^{def}	80 ^{abc}	57.5 ^{efgh}
	4	55 ^{def}	42.5 ^{cde}	90 ^{ab}	67.5 ^{de}
	8	77.5 ^{abcd}	60 ^{bc}	95 ^a	85 ^{bc}
	Mean	55.63 ^b	41.88 ^c	83.75 ^{bc}	63.75 ^c
F between treat. concn. and ages df	15	1.37	1.52	1.28	1.06
F between concentrations df	3	38.18		30.2	3
F between treatment df	3	75.08		21.31	3

Data presented in **table 1** indicate that the mortality percentage after treatment of the 2nd and 4th instar larvae of *S.littoralis* and *H. brunneipennis* increased gradually with increasing concentrations of all the insecticides.

The present results revealed that the tested Bt and fungus formulations had insecticidal activity against 2nd and 4th instar larvae of *S.littoralis* and *H. brunneipennis* larvae, where DipelDf highly killed the insect larvae both insect species, followed by W-Bus, Biofly and Protecto, respectively. These results agree with **Haggag, (2013)** who reported that DipelDf, Dipel 2x and Delfin highly killed *S.littoralis* larvae, followed by Agry, Protecto and Agerin, respectively. **Kaur (2000)**, also, reported that *B.thuringiensis* applied for controlling of lepidopteran, dipteran and coleopteran insects for decades. **Herrnstadt and Soares (1989)** reported

that *B.thuringiensis* 7.6x10⁷ spores/ml solution, caused 80% mortality against alfalfa weevil. The surviving weevil larvae were stunted and ceased feeding. Lower concentrations resulted in minimal levels of mortality, but caused significant levels of feeding inhibition, these inhibited larvae will not survive to adulthood in the field. *B.thuringiensis* produced more than 93% mortality on first instar larvae of *Spodoptera frugiperda* and *Peridroma saucia* (**Alvarez et al., 2009**). *B.thuringiensis* Berliner is a promising agent for microbial control of agriculturally and medically important insects (**Souza et al., 2009**). The difference in activity might be due to the presence or absence of biologically active Cry toxins, their relative amounts and additive/ synergistic effect of these toxins in the formulations. **Shelton et al., (1993)**. **Karthikeyan and Selvanarayanan (2011)** reported that the bioassay with *B. bassiana* against *S. litura*, percent

mortality increased from 33.33 to 86.67 as the dose was increased from 0.15 to 0.25 %.

Susceptibility test

Table (2) reveals the LC₅₀ values of the tested compounds against 2nd and 4th instar larvae of *S.littoralis* and *H. brunneipennis* recording 1.13&1.47; 2.75&6.47; 9.08& 14.90 and 2.015,5.05 gm/l, for DipelDf, W-Bus, Protecto and Biofly against 2nd and 4th instar larvae of *S.littoralis* , respectively while those were 0.84, 0.14; 0.59, 0.44; 1.94& 1.41 and 2.53& 5.20 for *H. brunneipennis*, respectively.

According to the LC₅₀ values, DipelDf was the highest toxic to *S.littoralis* and *H. brunneipennis* than the other 3 compounds. The toxicity values of DipelDf was significantly higher than that others.

Effects of LC₅₀ of Bt and *B.bassiana* formulations on pupation and adult emergence percentage.

The initial and residual effects of Bt and *B.bassiana* formulations at four time intervals (2, 4, 6, and 8 days) post application against 2nd and 4th instar larvae of *S.littoralis* and *H. brunneipennis* (at 6 days after treatment) are shown in **tables (3 and 4)** . Data in **table (3)** revealed that, treatment with all the tested compounds reduced pupation and adults emergence percentages and ,also, reduced the population of *S.littoralis* larvae compared to the control at initial and residual time intervals (2, 4, 6 and 8 days) to record 51, 43.6 , 35.9 and 38.5 at 2 time; 37.5 , 30 , 25 and 27.5 at 4 days; 30, 27.5,15 and 20 at 6 days and 25.6, 20.5,7.7 and 12.8 % larval mortalities at the 8 days, respectively on 2nd instar larvae and recored 42.5 , 37.5 , 30and 32.5 ; 30, 25, 20 and 20; 20, 17.5, 7.5 and 12.5; 17.5, 15, 5 and 10 at (2, 4, 6, and 8 days), respectively on 4th instar larvae.

The results in **Table (3)** indicated also that DipelDF and W-Bus decreased both pupation and adult

emergence percentages at (2, 4, 6, and 8 days) more than Biofly and Protecto compared to the control which recorded pupation and emergence rates of 97.5 and 100%, respectively.

The results of reduction percentage of *H. brunneipennis* population, pupation and adult emergence percentages after four indicating time intervals are summarized in **Table (4)**. Data showed that the mean percentage of cumulative larval mortality, pupation and adult emergence percentages of *H. brunneipennis* after four indicating time intervals of application(2, 4, 6, and 8days) varied among the all treatments and control. The reduction was 60 , 55, 40, and 45; 45, 37.5, 20 and 25% larval mortality on the 2nd instar larvae and 52.5, 47.5, 35 and 47.5; 27.5, 20, 10 and 15 % larval mortality on the 4th instar larvae at the initial time interval (2 day) and 6 days of application for DipelDf, W-Bus,Protecto and Biofly, respectively. These results agree with **El-Ghar et al.,(1995)** working with *Bacillus thuringiensis* and Abamectin against *S.littoralis* , with a pronounced decrease of pupation (36%) after Abamectin treatment. **Mohamed and Mahmoud, (2008)** reported that the rates of pupation and the emergence of moths of *S.littoralis* were reduced by all tested insecticides (Dipel 2x, Agrin, BioGuard, Biofly and Spinosad), respectively as compared to the control. *Beauveria bassiana* caused significant decrease in pupal survival with the malformation among *S.littoralis* pupae (**Emara and Hefnawy,2000**). Hyphomycete fungi cause fatal infection to the immature stages of *S.littoralis*, this may due to the disruption of normal metabolism, and damage of target tissues such as fat body or alter hormone balance (**Meshrif et al., 2007**).

From the above results and based on the LC₅₀ values, DipelDf, proved as the highest toxic to *S.littoralis* and *H. brunneipennis* than that of the other compounds, followed by ,W-Bus,Biofly and Protecto.

Table2. Lethal concentration of Bt and fungi formulations against *S. littoralis* and *H. brunneipennis* larvae

Formulation	<i>S. littoralis</i> gm/L (spores/ml)		<i>H. brunneipennis</i> gm/L (spores/ml)			
	2 nd	4 th	2 nd	4 th		
Dipel DF	LC ₅₀	1.13	1.47	LC ₅₀	0.14	0.84
	slope	3.07	2.34	slope	0.69	1.34
W-Bus	LC ₅₀	2.015	5.05	LC ₅₀	0.44	0.59
	slope	1.03	0.85	slope	0.39	1.48
Protecto	LC ₅₀	9.08	14.90	LC ₅₀	1.41	1.94
	slope	0.90	0.80	slope	1.12	1.24
Biofly	LC ₅₀	2.75	6.47	LC ₅₀	1.20	1.53
	slope	1.23	1.01	slope	1.40	1.63

Table 3. Initial and residual effect of the tested Bt and fungal formulations against 2nd and 4th instar larvae of *S. littoralis* at 6 days after treatment

Formulations	Accumulative larvae mortality % after indicated time intervals(days)											
	Initial kill			Residual effect								
	Accumulative 2 day			Accumulative 4 day			Accumulative 6 days			Accumulative 8 days		
	%Corrected mortality	%Pupation	%Adults emergence	%Corrected mortality	%Pupation	%Adults emergence	%Corrected mortality	%Pupation	%Adults emergence	%Corrected mortality	%Pupation	%Adults emergence
Dipel DF												
2 nd	51	47	58	37.5	62	64	30	70	68	25.6	72	76
4 th	42.5	58	61	30	70	71	20	80	75	17.5	82	78
W-Bus												
2 nd	43.6	55	63	30	70	68	27.5	72	72	20.5	78	77
4 th	37.5	62	68	25	75	76	17.5	82	79	15	85	79
Protecto												
2 nd	35.9	63	76	25	75	80	15	85	82	7.7	90	89
4 th	30	70	86	20	80	84	7.5	92	89	5	95	92
Biofly												
2 nd	38.5	60	71	27.5	73	76	20	80	84	12.8	85	88
4 th	32.5	67	81	20	80	81	12.5	87	83	10	90	91
Control												
2 nd	-	95	94.7	-	97	95	-	97.5	94.9	-	97.5	100
4 th	-	97.5	97.4	-	92.5	94.5	-	97.5	100	-	97.5	100

Table 4. Initial and residual effect of the tested Bt and *B.bassiana* formulations against 2nd and 4th instar larvae of *H. brunneipennis* at 6 days after treatment

Formulations	Accumulative larvae mortality after indicating time intervals(days)											
	Initial kill			Residual effect								
	Accumulative 2 day			Accumulative 4 day			Accumulative 6 days			Accumulative 8 days		
	%Corrected larval mortality	%Pupation	%Adults emergence	%Corrected larval mortality	%Pupation	%Adults emergence	%Corrected larval mortality	%Pupation	%Adults emergence	%Corrected larval mortality	%Pupation	%Adults emergence
Dipel DF												
2 nd	60	40	44	52.5	47	47	50	50	60	45	55	64
4 th	52.5	47	53	45	55	59	32.5	67	63	27.5	72	69
W-Bus												
2 nd	55	45	50	47.5	52	52	45	55	55	37.5	62	68
4 th	47.5	52	57	40	60	62	27.5	72	66	20	80	72
Protecto												
2 nd	40	60	56	40	60	58	27.5	73	69	20	80	78
4 th	35	65	65	27.5	72	69	20	80	75	10	90	81
Biofly												
2 nd	45	55	54	40	60	58	32.5	67	63	25	75	73
4 th	47.5	52	62	35	65	69	25	75	70	15	85	76
Control												
2 nd	-	95	97	-	92.5	97	-	95	100	-	92.5	100
4 th	-	95	100	-	97.5	92	-	97.5	100	-	97.5	100

References

- Abbott, W.S. (1925). A method for computing the effectiveness of an insecticide. *J. Econ. Entomol.*, 18(2) : 256-267.
- Adham, K. Fatma; Eman, M.Rashad; S.F. Ibrahim and Enas, E. Nasr, 2009. Host plants shifting affects the biology and biochemistry of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). *Egypt. Acad. J. Biolog. Sci.*, 2(1): 63-71.
- Ahmed, M. 2009. Observed potentiation between pyrethroid and organophosphorus insecticides for the management of *Spodoptera litura* (Lepidoptera: Noctuidae). *Crop Protection*, 28: 264-268.
- AL- Doghairi, M. A. and E. A. EL-Hag, 2003. Effect of several biopesticides on alfalfa weevil larvae, *Hypera brunneipennis* (Boheman). *Pakistan Journal of Biological Sciences*. 6 (8) 777- 781.
- Alvarez, A.; Virla, E. G.; Pera, L. M. and Baigori, M. D. 2009. Characterization of native *Bacillus thuringiensis* strains and selection of an isolate active against *Spodoptera frugiperda* and *Peridroma saucia*. *Biotechnol. Lett.*, 31: 1899-1903.
- Asi, M.R., M.H. Bashir, M. Afzal, B.S. Khan, M.A. Khan, M.D. Gogi, K. Zia and M. Arshad, 2012. Potential of entomopathogenic fungi against larvae and eggs of *Spodoptera litura* (Lepidoptera: Noctuidae). *Pak. Entomol.*, 34(2): 151-156.
- Bhattacharya, A.K.; Mondal, P.; Ramamurthy, V.V. and Srivastava, R.P. 2003. *Beauveria bassiana*, A potential bioagent for innovative integrated pest management programme. In: Srivastava, R.P. (Ed.), *Biopesticides and Bioagents in Integrated Pest Management of Agricultural Crops*. International Book Distributing Co. Lucknow 860pp.
- Dhir, B.C., H.K. Mohapatra and B. Senapati, 1992. Assessment of crop loss in groundnut due to tobacco caterpillar, *Spodoptera litura* (F.) *Indian J. Plant Prot.*, 20 (7- 10) : 215-217.
- Duncan, D. B. 1955. The Multiple Range and Multiple F- Test. *Biometrics* 11: 1 – 42.
- El- Ghar, G.E.S.A.; Radwan, H. A. S. ; El- Bermany, Z. A.; Zidan, L. T. M. 1995. Sublethal effects of avermectin B1, betaexotoxin of *Bacillus thuringiensis* and diflubenzuron against cotton leafworm (Lepidoptera: Noctuidae). *J. Appl. Entomol.*, 119: 309 – 313.
- El – Hawary, F. M. and Abd El- Salam, A. M. E. 2009. Laboratory bioassay of some entomopathogenic fungi on *Spodoptera littoralis* (Boisd.) and *Agrotis ipsilon* (Hufn.) larvae (Lepidoptera: Noctuidae) *Egypt. Acad. J. Biolog. Sci.*, 2(2) : 1 – 4.
- Emara, T. E. and Hefnawy, M. A. 2000. Biological activity of some fungal extract against the development of the cotton leafworm *Spodoptera littoralis* *J. Egypt. Ger. Soc. Zool.*, 33(E) : 217- 225.
- Finney, D. F. 1971. *Probit analysis* 3rd edition. Cambridge University, London. 318PP.
- Haggag, Karima, H. E. 2013. Changes in protein profile of cotton leafworm, *Spodoptera littoralis*, induced by Bt formulations stored at cold and hot storage conditions. *Nature and Science*, 11 (7): 77- 85.
- Hammad, S. M., S. El- Sherif, M. M. Hosny and A. I. El- Deeb. 1967. The biology of *Hypera brunneipennis* Boh. *Bull. Soc. Ent. Egypte. Li.* : 251 – 256.
- Herrnstadt, Corinna and George, G. Soares., 1989. Cotton boll weevil, alfalfa weevil, and corn rootworm via contact with a strain of *Bacillus thuringiensis*: 4, 797, 276.
- Kamel, Aida S. ; Mona, Ab – El Aziz F. and Nehad, El- Barky M.. 2010. Biochemical effects of three commercial formulations of *Bacillus thuringiensis* (Agerin, Dipel 2x and DipelDf) on *Spodoptera littoralis* larvae. *Egypt. Acad. J. Biolog. Sci.*, 3 (1) : 21- 29.
- Karima, H. E. Haggag, 2013. Changes in protein profile of cotton leafworm, *Spodoptera littoralis*, induced by Bt formulations stored at cold and hot storage conditions. *Nature and Science* 11 (7) 77- 85.
- Karthikeyan, A. and Selvanarayanan, V. 2011. In vitro efficacy of *Beauveria bassiana* (Bals.) Vuill. and *Verticillium lecanii* (Zimm) Viegas against selected insect pests of cotton. *Science and Technology*, 3 (2): 142- 143.
- Kaur, S. 2000. Molecular approaches towards development of novel *Bacillus thuringiensis* biopesticides. *World Journal of Microbiology and Biotechnology*, 16: 781- 793.
- Meshrif, W. S. ; Barakat, E. M. S. ; Rohlf, M. ; Shehata, M. G. ; Seif, A. I. and Hegazi, M. A., M. 2007. Evaluation of testing four hyphomycete fungi against the development of the cotton leafworm *Spodoptera littoralis* (Lepidoptera: Noctuidae). *J. Egypt. Acad. Soc. Environ. Develop.*, 8(3) : 11- 20.
- Mohammed A. Al- Doghairi and Eltayeb El Hag. 2003. Effect of several biopesticides on alfalfa weevil larvae, *Hypera brunneipennis* (Boheman). *Pakistan Journal of Biological Sciences*. 6 (8) 777- 781.
- Mohamed, E. M., Hanan, F. Abdel- Hafez and Mahasen A. Abdel Aziz. 2010. Effect of some chemical additives on the potency of *Bacillus thuringiensis* against the cotton leafworm *Spodoptera littoralis*. *Egypt. J. Agric. Res.*, 88(1), 103-112.
- Mohamed, A. M. and Mahmoud, F. M. 2008. Effect of biorational insecticides on some biological aspects of the Egyptian cotton leafworm

- Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae) . Plant Protect. Sci., 44: 147- 154.
- Prayogo, Y., W.Tengkano and D. Marwoto, 2005. Prospect of entomopathogenic fungus *Metarhizium anisopliae* to control *Spodoptera littoralis* on soybean. J. LitbangPertanian, 24(1) : 19-26.
- Sanchis V.; Gohar, M .; Chaufaux, J. ; Arantes, O. ; Meier, A. ; Agaisse, H. ; Cayley, J. and Lereclus, D. 1999. Development and field performance of a Broad- Spectrum Nonviable Asporogenic Recombinant Strain of *Bacillus thuringiensis* Greater Potency and UV Resistance. Applied Environmental Microbiology, 65 (9) : 4032- 4039.
- Shelton, A. M.; Robertson, J. L. ; Tang, J. D. ; Perez, C.;Eigenbrode, S. D.; Preisler, H. K.; Wilsey, W. T. and Cooley, R. J. 1993. Resistance of diamondback moth (Lepidoptera :Plutellidae) to *Bacillus thuringiensis* sub species in the “eld. Journal of Economic Entomology, 86: 697-705.
- Souza, Jr, J. D. A. de; Jain, S.; de Oliveira, C. M. F.; Ayres, C. F. and Lucena, W. A. 2009. Toxicity of *Bacillus thuringiensis* israeliensis – like strain against *Spodoptera frugiperda* .BioControl, 54: 467- 473.

الملخص العربى

تقييم لبعض المستحضرات التجارية على دودة ورق القطن وسوسة ورق البرسيم

نارمين أحمد يوسف¹ و عاطف عبد الجيد²

قسم وقاية النبات - كلية الزراعة - جامعة الفيوم - الفيوم - مصر .

تم تقييم فاعلية بعض المستحضرات البكتيرية (الدايل دى إف ، دايلو بص، بروتيكتو) والفطرية بيوفلاى ضد كل من العمر اليرقى الثانى والرابع لدودة ورق القطن و سوسة ورق البرسيم تحت الظروف المعملية. وأوضحت النتائج ان نسب الموت عند استخدام المستحضرات البكتيرية على كل من العمر اليرقى الثانى والرابع لدودة ورق القطن تراوحت بين 40 - 100 % ، 32.5 - 92.5 فى حين وصلت الى 100 ، 85 و 100 % على العمر اليرقى الثانى والرابع لسوسة ورق البرسيم على التوالي عنداستخدام أعلى تركيز بعد 7 ايام من المعاملة مقارنة ب 77.5 و 60 % على دودة ورق القطنو 95 ، 85 % لسوسة ورق البرسيم عند استخدام البيوفلاى. وبناء على قيم كان الدايل افضل المستحضرات المستخدمة ضد كل مندودة ورق القطن و سوسة ورق البرسيم مقارنة بالمستحضرات الاخرى. كما إنخفضت نسب خروج كل من العذارى والحشرات الكاملة عند المعاملة بالمستحضرات السابقة مقارنة بالكنترول.