

Physiochemical characteristics and therapeutic effects of lyophilized Bee venom and Royal jelly

Sharoba**, A.M.; EL-Tanahi, H.H.; Hafez, H.N.; Hendawy*, H.A. and Yousef, S.Y.A.

Food Science Department, Faculty of Agriculture, Benha University.

* Anesthesia, Intensive Care and Pain Relief Department, Faculty of Medicine, Suez Canal University

**Corresponding author: Dr. Ashraf Sharoba

Food Science Department, Faculty of Agriculture, Benha University, Qaluobia, Egypt.

E-mail: ashraf_sharoba@yahoo.com

Tel. 00201221463079 Fax: 0020132467786

Abstract

This study was investigating the chemical and physical analysis of bee venom and royal jelly before and after lyophilization process and found that the lyophilization process improved the efficiency of the enzymes and active ingredients as a result of low moisture to 5%. The use of bee venom and royal jelly improved the therapeutic efficiency of some diseases: improving the sexual efficiency, the functions of the kidney and decrease the rate of glucose by using the doses of 500 mg and 1 gram of (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey) / kg of body weight orally daily for adult male rats for a month. This study showed that the treatment of adult male rats by 1 gram (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey) / kg of body weight orally alone resulted in increasing the weights of the testis and the body of the epididymus, sperm count, testosterone hormone, the percentage of live sperm, and glutathione level, accompanied with decrease in malondialdehyde level and the percentage of sperm abnormality. The oral administration of (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey) in doses of 500 mg and 1 g/kg body weight today for one month increased urea concentration in serum and decreased in urine, urea clearance in urine was also decreased as compared to control group. (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey) in a dose of 1 g/kg body weight daily decreased serum, urine and creatinine levels. Decrease in serum glucose level was also observed when the test animals were treated with (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey) in a dose of 500mg and 1g /kg body weight daily. Remarkable for one month was showed increasing in body weight of adult rats that treated with 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey) /kg of body weight compared with control group. This study recommended that using mixtures of lyophilized bee venom and royal jelly which are dissolved in honey as therapy of sexual efficiency, kidney and sugar diseases.

Keywords: Bee venom, royal jelly, sexual efficiency, therapeutic efficiency.

Introduction:

*Eros, stung by a bee,
ran away and cried for plea:
Venus, mother, I cry,
please help me or I'll die
What a terrible disgrace –
a dragon bit me on my face
Venus comforting her son
Speaking with a mocking fun -
The little bee's tiny sting
Is for you an earnest thing
But more painful and real hard
are your stings in human's heart
Anacreontean songs, 6 BC*



Venus, Eros and the bees, by A. Dürer, 1514

It was difficult for Eros to bear the bee stings which like his arrows and both painful and healing.... . Already in the early ancient civilizations knew about the healing found virtues in the painful

bee stings. Bee stings are probably one of the first natural cures for arthritis. In the ancient civilization of China, India, Egypt, Babylon and Greece bee venom was used for apitherapy (Urtubey, 2005).

Whether the humans began keeping bees because of the healing effects of their stings or to get honey, or for both reasons, we do not know (<http://www.bee-hexagon.net> (2012)). Also, the use of honey and other bee products in human treatments traced back thousands of years and healing properties are included in many religious texts including the Veda, Bible and Quran (El-Banby, 1994). Apitherapy is the use of honey bee products for medical purposes, this include bee venom, raw honey, royal jelly, pollen, propolis and beeswax. Whereas bee venom therapy is the use of live bee stings (or inject able venom) to treat various diseases such as arthritis (Beck, 1999), rheumatoid arthritis, multiple sclerosis (MS), lupus, sciatica, low back pain, and tennis elbow to name a few. It refers to any use of venom to assist the body in healing itself. Bee venom contains at least 18 pharmacologically active components including various enzymes, peptides and amines (Habermann and Jentsch, 1999).

Sulfur is believed to be the main element in inducing the release of cortisol from the adrenal glands and in protecting the body from infections. Contact with bee venom produces a complex cascade of reactions in the human body. The bee venom is safe for human treatments, the median lethal dose (LD₅₀) for an adult human is 2.8 mg of venom per kg of body weight (Crane, 1990), for example a person weighing 60 kg has a 50% chance of surviving injections totaling 168 mg of bee venom (Tumanov and Osipova, 1996). Assuming each bee injects all its venom and no stings are quickly removed at a maximum of 0.3 mg venom per sting, 560 stings could well be lethal for such a person. For a child weighing 10 kg, as little as 93.33 stings could be fatal. However, most human deaths result from one or few bee stings due to allergic reactions, heart failure or suffocation from swelling around the neck or the mouth. Compared with other human diseases, accidents and other unusual cases, the bee venom is very safe for human treatments (Bogdanov, 2011a).

Bee venom: Honeybee venom is produced by two glands associated with the sting apparatus of worker bees. Its production increases during the first two weeks of the adult worker's life and reaches a maximum when the worker bee becomes involved in hive defenses and foraging. It diminishes as the bee gets diminishes as the bee gets older.

Royal jelly: is a thick, extremely nutrition, milky white, creamy liquid secreted by the hypo pharyngeal glands of worker bees (*Apis mellifera*) in relation to sexual determination of the bee (Antinelli et al., 2003). Considered as the major cause for difference between queen and bee workers, royal jelly is appreciated as a dietary complement because of its composition. Royal jelly is an essential food for the queen bee larvae and the queen herself. All larvae fed royal jelly for three days, but the queen bee eats royal jelly exclusively which makes her fertile and able to live to seven years. Queen bees will produce

2000 eggs per day, with each day brood equal to 2.5 times her body weight (Leung et al., 1997). In contrast, worker bees are sterile and live just seven to eight weeks. Royal jelly contains considerable amounts of proteins, amino acids including 8 essential amino acids, hormone rich substance (testosterone has been identified in extremely small quantities in royal jelly about 0.012g/g fresh weight <http://www.goldinnature.com/apitherapylinks.html> (2004), royal jelly also contains lipid, sugars, vitamin A, C, D & E, minerals are in descending order: (K, Ca, Na, Zn, Fe, Cu and Mn), enzymes antibiotic components and an abundance of nucleic acid-DNA and RNA.

Gelatin, one of the precursors of collagen, is also found in royal jelly, collagen is a powerful anti-aging element that helps preserve the youth of the body (Compston, 2001). Gelatin is known to have several diverse physiological and pharmacological functions, these include vasodilative, hypotensive, ant hypercholesterolemia, and anti-tumor activities (Narita et al. 2006). Royal jelly has been found to be of great help in boosting the body resistance to the harmful side effect of chemotherapy and radiotherapy <http://www.apitherapy.org> (2001). Also, contains gamma globulin, which helps the immune system to fight infections. It also contains sterols, phosphorous compounds and acetylcholine, which are needed to transmit nerve messages from cell to cell.

Al-Taei (2003) demonstrated that, the reactive oxygen species produced by administration of hydrogen peroxide are responsible for the pathophysiological changes of the male reproductive system and induced defect in the histophysiological aspect of this system in rats.

Polyunsaturated fatty acids and phospholipids are key constituents in the sperm cell membrane and are highly susceptible to oxidative damage. Sperm produce controlled concentrations of reactive oxygen species, such as the superoxide anion, hydrogen peroxide, and nitric oxide, which are needed for fertilization; however, high concentrations of these free radicals can directly damage sperm cells (Ebisch et al., 2006). On another hand, the active ingredients in bee venom and royal jelly are active due to lyophilization, Bogdanov (2011b).

Honey: Honey is the natural sweets substance produced by honeybees from the nectar of blossoms or from the secretion of living parts of plants or excretions of plant sucking insects on the living parts of plants, which honeybees collect, transform and combine with specific substances of their own, store and leave in the honey comb to ripen and mature. This is the general definition of honey in the Codex Alimentarius (1989).

The current study was aimed to investigate effect the administration different concentrations 1g, 500 mg of (1mg bee venom +500mg royal jelly +25mg honey) orally for one month daily to the male adult

rats induced oxidative stress by hydrogen peroxide on sexual efficacy and glutathione, malondialdehyde levels, and to study the effect of this administration on the glucose level and serum in rats.

Materials and methods:

Materials

Bee venom, royal jelly and honey were obtained as pure substance from GHADA Company – Borgalarb- Alexandria-Egypt, in January, February, April, June, September, October and December 2013.

Bee venom extraction:

The venom collection was conducted by Moshtohor device according to **Khattab (1997)**.

Lyophilization of bee venom:

Add 1 ml of distilled water per 1 gram of raw bee venom and filtering by nomination paper 0.02 to get rid of impurities. Then the venom was freezing at -50°C for 24 hours and put in lyophilization device (invented by the researcher), at -10°C and a pressure at -0.8 bar to reach to sublimation process for 48 hours, finally, getting lyophilized bee venom as a powder with a suitable moisture content 6.0 - 8.0%.

Lyophilization of royal jelly:

Add 5 ml of 13% ethanol per 1 gram of raw royal jelly then freezing at -70°C for 24 hours. After that, put royal jelly in lyophilization device (invented by the researcher), at -10°C and a pressure at -0.8 bar to reach to sublimation process for 48 hours. Finally, getting a lyophilized royal jelly as a powder with suitable moisture content 3.41- 6.62%. Then store in dark jars at room temperature (25°C - 30°C). After that dissolve 1mg of lyophilized bee venom + 500 mg of lyophilized royal jelly in 25 mg bee honey, according to **DGHCP (2002)**.

Methods:

Chemical analysis:

Moisture, crude protein, lipid and ash contents were determined according to the methods described in **AOAC (2005)**.

Fructose, glucose, sucrose, 10-hydroxy-2deconic acid, melitine, phospholipids A and hyaluronidase were determined according to the method by H P L C analysis.

pH value of device model MA 5736, Metrel, Iskra, Slovenia was determined according to the method described by **AOAC (2005)**.

Animal Experimental:

Sixty three adult male albino rats were obtained from the animal house of the Medical Research Center, Alexandria, was aged 2-3 month and weights

ranged from 220 - 280g. They were housed in polypropylene cages under controlled condition of temperature ($24-26^{\circ}\text{C}$) and lighting (12hours light/12hours dark). The rats were supplied with standard diet and tap water.

The adult male rats were randomly divided into six groups. The first three groups were 12 rats/group and the second three groups were 9 rats/ group. The first group received tap water serve as control. The second group received 0.5% hydrogen peroxide (H_2O_2) daily in drinking water for one month to be compared with the sample (**Abdul-Rahman, 1995**). The third group received 0.5% (H_2O_2) daily in drinking water for one month concomitant with (1mg of lyophilized bee venom+500mg of lyophilized royal jelly +25mg honey) at 1g/kg body weight dissolved in distilled water and given daily for one month orally by gavages needle (**Mishima et al ., 2005**). The fourth, fifth and sixth groups received (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey) at a dose 1g/kg body weight orally daily. The weight of rats recorded weekly. At the end of experiment blood samples were collected into clean dry centrifuge tubes allowed to clot, serum separated after centrifugation at 1500 rpm for 15 minute for testosterone hormone assay, using Enzyme Linked Immune sorbent Assay (ELISA) (BioCheck Company, USA). Rats were sacrificed by ether administration. The abdominal cavity was then opened; the weight of testis, epididymal, seminal vesicles and prostate were recorded. The test is placed in ice normal saline for glutathione estimation using Moron method as described by malondialdehyde (MDA) estimation using Gilbert method. The epididymis was dissected out, sectioned and immediately the content of the tail of each epididymis was squeezed gently in clean watch glass diluted 10 times with isotonic solution of sodium citrate 2.9% at 37°C , take one drop from isotonic solution on slide and added one drop of eosin - nigrosin stain and made smear, this technique was used for the percentage of live/dead and for morphological abnormal sperms to be counted (**Al-Sadi, 2001**). The content of the head of epididymis was squeezed immediately in clean watch glass contained 9.8 ml buffer formalin with 0.1 ml eosin 5%, this was used for counting the sperm concentration using hemocytometric technique (**Bearden et al., 2004**), and comparing with the control group, (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey) in dose of 50 and 100 mg/kg body weight daily which obtained on it the level of creatinine in serum and urine, finally we obtained glucose level in serum and urine to rats which administrated by 50 and 100 mg/kg body weight of (1mg of lyophilized bee venom +500mg of lyophilized royal jelly +25mg honey).

Statistical analysis:

Data were analyzed statistically using one way analysis of variance as determined by **Petrie and Watson (1999)**.

Results and discussion:

a. Effect of Lyophilization on the chemical characteristic of royal jelly and bee venom:

a.1. Chemical characteristic of royal jelly

According to table (1) we can recognized that the reduction in moisture content of royal jelly samples was ranged between 92 – 95% after lyophilized the samples. This was in accordance with previous scientific researches (**Messia et al., 2005 and Bogdanov (2011a)**).

The proportion of the proteins in royal jelly was large. From Table (1), the average value of total protein was increased from 8.62 – 9.14 % to 25.90 – 27.44% after lyophilization. Similar results were reported by **Simúth (2001) and Kim et al., (2010)**.

The same trend was observed with royal jelly total lipids in Table (1). There was increasing in the total lipids average 5.19 – 5.72 % to 8.40 – 9.10 % after lyophilization. This is in accordance with **Sabatini et al., (2009) and Kodai et al.,(2007)**.

The concentration of the most abundant sugars fructose was ranged between 7.92 – 8.11% and after lyophilization ranged between 5.11 – 16.39%, glucose was 5.53 – 5.81% after lyophilization ranged from 10.21 – 10.98% and sucrose ranged from 0.52 – 0.61% and after lyophilization ranged between 1.20 – 1.32% (see Table 1). These values are comparable with the sugar levels reported in the literature (**Simúth, 2001 and Sabatini et al., 2009**).

There were some variations in values for individual samples which was normal for organic products.

According to **Simúth (2001)**, the average level of glucose in lyophilized royal jelly (10.21 – 10.98%) was higher than that of fructose (5.11 – 16.39%). **Sabatini et al., (2009)** reported that fructose was prevalent to glucose which was also confirmed in our study. Sucrose is always present but in highly variable concentrations. Thus the amount of analyzed royal jelly samples was insufficient to draw conclusions about the prevalence of glucose or fructose.

Royal jelly was highly acidic with pH 3.4 – 3.90 from Table (1), after lyophilization pH slightly decreased and became 3.29 – 3.81. The results were in accordance with **Lercker (2003) and Scarselli et al.,(2005)**.

Most of the organic acids are free with rather unusual structure rarely encountered in nature, mono- and dihydroxy acids and dicarboxylic acids with 8 and 10 carbon atoms (**Lercker et al., 1993**), the main acid being 10-hydroxy-2deconoic acid, which was antibacterial and immune activating (**Bachanova et al., 2002**), immune-modulating, anti-cancer (**Dzopalic et al., 2011**), anti-diabetes (**Okuda et al.,1998**), collagen promoting and skin protecting , anti-ulcer (**Fang,1994**), facilitates differentiation of brain cells (**Hattori et al., 2007**) antidepressant in mice experiments (**Ito et al., 2012**), promotes endothelial health, antihypertensive, antihyperlipidemia (**Izuta et al., 2009**) estrogenic(**Matsui et al., 2002**) anti-rheumatic and activation of TRPA1 and TRPV1 (induces thermogenesis and energy expenditure enhancement) (**Terada et al., 2011**).

HAD was ranged between 2.71 – 2.91% and after lyophilization it became 8.13 – 8.70% according to (**Bogdanov (2011a)**).

Table 1. Chemical composition of fresh and lyophilized royal jelly (Mean ± SE).

Parameters	Royal Jelly							
	(February)		(April)		(June)		(September)	
	A	B	A	B	A	B	A	B
Moisture%	66.28 ± 0.01	5.27 ± 0.01	64.17 ± 0.01	3.41 ± 1.66	67.21 ± 0.01	5.11 ± 0.01	68.15 ± 0.01	5.62 ± 0.01
PH	3.87 ± 0.01	3.62 ± 0.01	3.90 ± 0.06	3.81 ± 0.01	3.60 ± 0.06	3.50 ± 0.06	3.40 ± 0.06	3.29 ± 0.01
Protein %	9.14 ± 0.03	27.44 ± 0.01	8.92 ± 0.01	26.41 ± 0.01	8.62 ± 0.01	25.90 ± 0.06	8.91 ± 0.01	26.91 ± 0.01
Fructose %	8.11 ± 0.01	16.39 ± 0.01	7.92 ± 0.01	15.11 ± 0.01	7.95 ± 0.01	16.12 ± 0.01	7.99 ± 0.01	16.11 ± 0.01
Glucose %	5.53 ± 0.01	10.21 ± 0.01	5.71 ± 0.01	10.61 ± 0.01	5.62 ± 0.01	10.41 ± 0.01	5.81 ± 0.01	10.98 ± 0.01
Sucrose %	0.55 ± 0.01	1.21 ± 0.01	0.52 ± 0.01	1.20 ± 0.01	0.61 ± 0.01	1.32 ± 0.01	0.58 ± 0.01	1.28 ± 0.01
Total sugar%	14.19±0.01	27.81±0.01	14.15±0.01	26.92±0.01	14.18±0.01	27.85±0.01	14.38±0.01	28.37±0.01
Ash%	0.71± 0.01	2.51± 0.01	0.62± 0.01	2.12± 0.01	0.69± 0.01	2.41± 0.01	0.65± 0.01	2.45± 0.01
10-hydroxy-2deconoic acid %	2.71± 0.01	8.13± 0.01	2.86± 0.06	8.38± 0.01	2.91± 0.06	8.70± 0.06	2.82± 0.06	8.56± 0.01
Lipids %	5.72 ± 0.01	9.10 ± 0.06	5.59 ± 0.04	8.92 ± 0.01	5.32 ± 0.01	8.40 ± 0.06	5.19 ± 0.01	8.60 ± 0.06

A: Fresh royal Jelly.

B: Lyophilized royal Jelly.

a.2 Chemical characteristics of bee venom

Bee venom is a transparent liquid dries up easily even at room temperature, odorless, ornamental

pungent smell, a bitter taste, hydrolytic blend of proteins with basic pH (4.5 – 5.5) that is used by bees for defense (Schmidt and Buchmann (1999)). When coming into contact with mucous membranes or eyes, it causes considerable burning and irritation. Bee venom is soluble in water and insoluble in alcohol and ammonium sulfate. When it comes in contact with air it forms grayish-white crystals. Lyophilized venom takes on a light yellow color and some commercial preparations are brown, thought to be due to oxidation of some of the venom proteins. Bee venom contains a number of very volatile compounds which are easily lost during collection, it is considered a rich source of enzymes, peptides and biogenic amines, and it is specific weight (1.1331). The venoms of most stinging insects

including honey bees consisted of enzymes, protein, peptides, and a variety of smaller molecules. The pharmacological and biochemical activities of the various stinging insect venoms remarkably convergent. Most venom induces immediate pain. From the results in Table (2), the phospholipids was ranged from 1.20 – 1.50% and became 9.60 – 11.20% after lyophilization, hyaluronidase was ranged from 0.28 – 0.32% after lyophilization it became 2.0 – 2.20% and is capable of destroying red blood cells. Most hymenopterous venoms also contain low molecular weight peptides (Norman *et al.*, (2011)). On the otherhand lyophilized bee venom content of melitine was ranged between 52.0 – 56.0% and in accordance with Bogdanov (2011b).

Table 2. Chemical composition of fresh and lyophilized bee venom (Mean \pm SE).

Parameters	Bee Venom							
	(January)		(April)		(October)		(December)	
	A	B	A	B	A	B	A	B
Moisture%	90.0 \pm 0.58	8.0 \pm 0.06	86.0 \pm 0.58	6.0 \pm 0.06	86.0 \pm 0.58	6.0 \pm 0.06	87.0 \pm 0.58	6.20 \pm 0.06
Melitine %	6.20 \pm 0.06	52.0 \pm 0.58	7.50 \pm 0.06	55.0 \pm 0.58	7.50 \pm 0.06	55.0 \pm 0.58	7.60 \pm 0.06	56.0 \pm 0.58
Phospholipids A %	1.20 \pm 0.06	9.60 \pm 0.06	1.50 \pm 0.06	10.98 \pm 0.01	1.50 \pm 0.06	11.20 \pm 0.06	1.50 \pm 0.06	11.10 \pm 0.06
Hyaluronidase %	0.28 \pm 0.01	2.0 \pm 0.06	0.31 \pm 0.01	2.10 \pm 0.06	0.32 \pm 0.01	2.20 \pm 0.06	0.30 \pm 0.01	2.10 \pm 0.06
Ash %	0.06 \pm 0.01	0.46 \pm 0.01	0.07 \pm 0.01	0.50 \pm 0.01	0.07 \pm 0.01	0.51 \pm 0.01	0.07 \pm 0.01	0.51 \pm 0.01
Total sugar %	0.31 \pm 0.01	1.90 \pm 0.06	0.36 \pm 0.01	2.10 \pm 0.06	0.34 \pm 0.01	2.0 \pm 0.06	0.34 \pm 0.01	2.0 \pm 0.06

A: Fresh bee venom.

B: Lyophilized bee venom.

b. Effect of treatment by royal jelly and bee venom on body weight:

From the data in Table (3) demonstrated there was no differences between groups in the body weight after one, two and three weeks of treatment.

After three weeks of treatment the body weight was increased compared with the group which was treated with H₂O₂ and the control group. Antinelli *et al* (2003).

Table 3. Effect the treatment of 1g (1mg of lyophilized bee venom + 500mg of lyophilized royal jelly +25ml honey)/kg of body weight orally on body weights in rats receiving hydrogen peroxide for one month (Mean \pm SE).

Treated animals	Weight (zero time)	Weight after one week	Weight after two weeks	Weight after three weeks
Control	270.5 \pm 10.64	246.2 \pm 11.91	273.5 \pm 24.4	299.5 \pm 10.1
1	228.5 \pm 20.34	228.2 \pm 19.70	238.5 \pm 24.6	287.2 \pm 21.6
2	263.0 \pm 8.09	223.5 \pm 20.01	235.2 \pm 21.9	286.2 \pm 14.3
3	265.0 \pm 23.47	250.2 \pm 17.06	247.0 \pm 26.5	305 \pm 34.07

1: Hydrogen peroxide (0.5%) in drinking water for one month.

2: Hydrogen peroxide (0.5%) in drinking water for one month) +1g (1mg of lyophilized bee venom + 500mg of lyophilized royal jelly +25ml honey) /kg of body weight orally for one month.

3: 1g (1mg of lyophilized bee venom + 500mg of lyophilized royal jelly +25ml honey)/kg of body weight orally for one month.

C. Effect of treatment by royal jelly and bee venom on increase fertility and efficiency nationality:

Data in Table (4) showed that, the administration by hydrogen peroxide (0.5%) in drinking water for

one month did not affect the weight of testis, epididymis(head, body, and tail), prostate and seminal vesicles compared with control group value, while the administration by hydrogen peroxide (0.5%) in drinking water for (1 month) +1g (1mg of

lyophilized bee venom + 500mg of lyophilized royal jelly +25ml honey) /kg of body weight orally for one month were accompanied by increasing in the weight of testis and body of epididymis, whereas no changes in the weight of head and tail of epididymus, prostate and seminal vesicles compared with hydrogen peroxide group. On the other hand, the administration by (1mg of lyophilized bee venom +

500mg of lyophilized royal jelly + 25ml honey) /kg of body weight orally for one month increased the weight of testis and body of epididymis, whereas no changes in the weight of head and tail of epididymus, prostate and seminal vesicles compared with hydrogen peroxide group, and this results were in agreement with those obtained by **Strum et al. (2002) and Abreu et al. (2000)**.

Table 4. Effect the treatment of 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally on the weight of the testis, epididymis (head, body, tail) prostate and seminal vesicle in rats receiving hydrogen peroxide for one month (Mean ± SE).

Treated animals	Testis mg/100g B.Wt.	Head of epididymus mg/100g B.Wt	Body of epididymus mg/100g B.Wt	Tail of epididymus mg/100g B.Wt	Prostate mg/100g B.Wt	Seminal vesicle mg/100g B.Wt
Control	473.4±27.7	79.2±4.2	21.1±0.9	93.2±7.4	447.5±44.2	100.9±5.9
1	501.5±10.8	85.0±3.9	20.8±0.6	92.9±5	431.35±35.3	108.9±7.4
2	604.9±212	89.4±5.5	23.7±0.3	93.4±2.9	455.3±33.2	99.9±6
3	636.1±21.7	76.5±4.1	24.1±0.3	90.4±1.5	427.1±26.7	105.7±6.4

1: Hydrogen peroxide (0.5%) in drinking water for one month.

2: Hydrogen peroxide (0.5%) in drinking water for one month +1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) /kg of body weight orally for one month.

3: 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally for one month.

Data in Table (5) that revealed that there was decrease in the sperm count in hydrogen peroxide group compared with control group, while, treatment with 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally and hydrogen peroxide caused increase in sperm count compared with hydrogen peroxide group and approximately returned to the normal control value(1.43). Data in the same Table showed increase in the sperm count by 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally treated group compared with hydrogen peroxide group, and 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally concomitantly with hydrogen peroxide group. Also, data in the same Table demonstrated that there was decrease in glutathione level in

hydrogen peroxide group compared with control group, while administration of 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) /kg of body weight orally with or without hydrogen peroxide caused increase in the glutathione level compared with hydrogen peroxide group. On the other hand, data in Table (5) showed that increase in malondialdehyde level in hydrogen peroxide group compared with control group while treatment with 1g(1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) /kg of body weight orally concomitant with hydrogen peroxide were no affect in the malondialdehyde level, but the treatment with 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally alone caused decrease in malondialdehyde level, **WHO (2002) and Krell (1996)**.

Table 5. Effect of the treatment of 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally on sperm count, glutathione, and malonaldehyde levels in the rats receiving hydrogen peroxide for one month (Mean ± SE).

Treated animals	Sperm concentration x10 ⁶	Glutathione µm/g	Malonaldehyde nm/g
Control	1.4320±0.02	1.04±0.02	264.82±12.48
1	0.80±0.02	0.59±0.02	311.0±17.6
2	1.3440±0.13	1.23±0.14	233.9±14.3
3	1.6260±0.02	1.26±0.02	219.7±5.7

1: Hydrogen peroxide (0.5%) in drinking water for one month.

2: Hydrogen peroxide (0.5%) in drinking water for one month + (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)(1g/kg orally) one month.

3: (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) (1g/kg orally) one month.

Data in Table (6) demonstrated that there was decrease in the percentage of the live sperms in

hydrogen peroxide group (84) compared with control group (91.6). Treatment with 1g (1mg of lyophilized

bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally concomitantly with hydrogen peroxide did not affect in the percentage of the live sperms compared with hydrogen peroxide group, Whereas the treatment with 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally alone caused increase in the percentage of the live sperms and returned to normal control value. Data in the same Table revealed that increasing in the percentage of sperms deformity in hydrogen peroxide group compared with control group. Also treatment with 1g (1mg of lyophilized bee venom +500mg of

lyophilized royal jelly+25ml honey)/kg of body weight orally with or without hydrogen peroxide caused decrease in the percentage of sperm deformity compared with hydrogen peroxide group. Data in the same Table showed decreasing in the testosterone hormone level in hydrogen peroxide group compared with control group. While, administration by 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey)/kg of body weight orally concomitant with or without hydrogen peroxide caused increase in testosterone hormone compared with hydrogen peroxide group, (Strum *et al.* 2002).

Table 6. Effect the treatment of 1g (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) /kg of body weight orally on the percentage number of live sperm, sperm deformity, and testosterone hormone concentration in rats receiving hydrogen peroxide for one month (Mean ± SE)..

Treated animals	Live Sperm%	Sperm Deformity%	Testosterone hormone mg/ml
Control	91.6±1.5	4.2±0.37	2.37±0.16
1	84±1.51	11.2±1.06	1.72±0.30
2	87±0.54	9.0±7.03	2.51±0.13
3	94.6±0.81	4.6±0.4	4.24±0.27

- 1: Hydrogen peroxide (0.5%) in drinking water for one month.
- 2: Hydrogen peroxide (0.5%) in drinking water for one month + (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) (1g/kg orally) one month.
- 3: (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) (1g/kg orally) one month.

d. Effect of treatment by royal jelly and bee venom in Improve kidney function:

Data in Table (7&8) show that oral administration with (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) orally in doses of 50 mg and 100 mg/kg body weight increased urea concentration in serum and decreased in urine. Urea clearance in urine was also decreased

as compared to control group. While, oral administration by (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) orally in a dose of 100 mg/kg body weight decreased serum and urine creatinine level which might be attributed to the increased cortisol level. (Abreu *et al.* 2000).

Table 7. Effect of oral administration of 50 and 100mg/kg body weight of (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) for 5 days on urea concentrations (mg/100ml) in serum and urine and urea clearance (ml/min) of rats (n=6)

Groups	Group (1) Control		Group (2) (50m/g kg b. wt)		Group (3) (100 m/g kg b. wt)				
	Urea concentration (mg %)		Urea concentration (mg %)		Urea concentration (mg %)				
	Serum	urine	Serum	Urine	Serum	Urine			
Values	40.19±0.93	145.00±5.69	0.019±0.0007	78.33±1.60	105.0±5.54	0.009±0.0004	88.28±1.28	46.33±3.13	0.0044±0.0003

* Mean ± SE

Table 8. Effect of oral administration of 50 and 100mg/kg body weight of (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) for 5 days on creatinine level (mg %) in serum and urine and creatinine clearance (ml/min) of rats (n=6)

Groups	Group (1) Control		Group (2) (50m/g kg b. wt)		Group (3) (100 m/g kg b. wt)	
	Creatinine concentration (mg %)		Creatinine concentration (mg %)		Creatinine concentration (mg %)	
	Serum	urine	Serum	Urine	Serum	Urine
		Greatinine clearance (mg %)		Greatinie clearance (mg %)		Greatinie clearance (mg %)

Values	0.29 ± 0.015	133.16 ± 4.00	2.57 ± 0.0007	0.30 ± 0.023	120.68 ± 3.98	2.89 ± 0.148	0.22 ± 0.019	88.75 ± 3.86	0.0855 ± 0.050
--------	--------------------	---------------------	---------------------	--------------------	---------------------	--------------------	--------------------	--------------------	----------------------

* Mean ± SE

e. Effect of treatment by royal jelly and bee venom in glucose concentrations in serum and urine of rats (Diabetics Rats):

Decreasing in serum glucose level was also observed when the test animals were treated with

(1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) orally in a dose of 50 and 100 mg/kg body weight (Table 9). Present findings suggest that (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) possessed insulin like effects (Munstedt *et al.* 2009).

Table 9. effect of oral administration of 50 and 100mg/kg body weight of (1mg of lyophilized bee venom +500mg of lyophilized royal jelly+25ml honey) for 5 days on glucose concentrations (mg/100 ml) in serum and urine of rats (n=6).

Rats Group No	Group (1) Control		Group (2) (50 mg/kg .b. wt)		Group (3) (100 mg/kg .b. wt)	
	Serum	Urine	Serum	Urine	Serum	Urine
Values	96.73* ± 0.584	0.00	88.42 ± 0.766	0.00	84.14 ± 1.21	0.00

* Mean ± SE

Reference

A.O.A.C. (2005) Official Methods of Analysis.

Association of Official Analytical, Chemists 18th ed., Washington, DC, USA.

Abdul-Rahman S.Y. (1995) Effect of starvation and experimental diabetes mellitus on glutathione and lipid peroxidation in tissues rats. Doctor's dissertation. College of Veterinary Medicine, University of Mosul, Iraq.

Abreu R.M.M.; Silva de Moraes R.L.M. and Malaspina O. (2000) Histological aspects and protein content of *Apis mellifera* L. worker venom glands: the effect of electrical shocks in summer and winter. *Journal Venom Animal. Toxins.*; **6**, 87 – 98.

Al-sadi A.A. (2001) Fertility and Artificial Insemination. 2nd ed College of Veterinary Medicine, University of Mosul. Iraq.

Al-Taei A.Y.J. (2003) Effect of vitamin C on some testicular function in rats exposed to oxidative stress induced by hydrogen peroxide. MSc. Thesis College of Veterinary Medicine, University of Mosul. Iraq.

American Apitherapy society. (2001). 5390 Grande road, Hillsboro, 45133. (937) 364-1108. <http://www.Apitherapy.Org/>. Gale Encyclopedia of Alternative Medicine. Gale Group.

Antinelli J.; Zeggane S.; Davico R.; Rognone C.; Faucon J. and Lizzani L. (2003) Evaluation of (E)-10-hydroxydec-2-enoic acid as a freshness parameter for royal jelly. *Food Chem.*; **80**:85-89.

Bachanova K.; Kludiny J.; Kopernicky J. and Simuth J. (2002) Identification of honeybee peptide active against *Paenibacillus* larvae larvae

through bacterial growth-inhibition assay on polyacrylamide gel. *Apidologie* **33** (3): 259-269.

Bearden H.J.; Fuguany T.W. and Willard S.T. (2004) Applied animal reproduction. 6th ed Mississippi State University.

Beck B.F. (1999) "Bee Venom Therapy. New York: Appleton-Century 1935; by Schmidt and Buchmann" "In. The Hive and the Honey Bee, Edited by Joe M. Graham, Dadant and Sons, Hamilton, Illinois".

Bogdanov S. (2011_b). Bee Venom: Composition, Health, Medicine : A Review . *Bee product science* 1-16. (<http://www.bee-hexagon.net>, 15 January 2012.)

Bogdanov S. (2011_a). Royal Jelly, bee brood: composition, health, medicine: A review. *Bee product science* 1-31. (<http://www.bee-hexagon.net>, 15 January 2012.)

Codex Alimentarius Commission (1989). Codex standards for sugars (honey). Supplement to codex Alimentarius volume III. Food and Agriculture Organization of the United Nations and World Health Organization, Rome.

Compston J.E. (2001) Sex steroids and bone. *Physiol Review*. **8**:419-447.

Crane E. (1990) "Bees and beekeeping": Science, Practice and World Resources. Cornstock Publication., Ithaca, NY., USA.; 593 pp,

D.G.H.C.P.: Directorate General Health and Consumer Protection (2002) Implementation of Regulation (EC) No 258/97 of the European parliament and the council of 27th January 1997 concerning novel foods and novel food ingredients. (SANCO D4) European Commission.

Dzopalic T.; Vucevic D.; Tomic S.; Djokic J.; Chinou I. and Colic, M (2011) 10-Dihydroxy-

- decanoic acid, isolated from royal jelly, stimulates Th1 polarising capability of human monocyte-derived dendritic cells. *Food Chemistry* 126 (3): 1211-1217.
- Ebisch I.M.W.; Pierik F.H.; Jong F.H.; Thomas C.M.G. and Steeger-Theunissen R.P.M. (2006)** Does folic acid and zinc sulphate intervention affect endocrine parameters and sperm characteristics in men. *International Journal Androl.*;29(2):339-345.
- El-Banby M.A. (1994)** "Honeybees in the Koran and in medicine. Al-Ahram Centre for Translation and Publication, Cairo, Egypt; 268 pp, (In Arabic). [6] Health and the Honey.
- Fang E.; Zhou. H.; Xu. H. and Xing. M. (1994)** Antiulcer effects of 10-hydroxy-2-decenoic acid in rats. *ZhongguoYaolixue Tongbao* 10 (2): 9-42.
- Habermann E. and Jentsch J. (1999)** "Sequenzanalyse des Melittins aus den tryptischen und peptischen Spaltstücken". *Hoppe-Seyler's Z. Physiol. Chem.* 348: 37-50, 1967."In. *The Hive and the Honey Bee*, Edited by Joe M. Graham, Dadant and Sons, Hamilton, Illinois, ").
- Hattori N.; Nomoto H.; Fukumitsu H.; Mishima S. and Furukawa S. (2007)** Royal jelly and its unique fatty acid, 10-hydroxy-trans-2-decenoic acid, promote neurogenesis by neural stem/progenitor cells in vitro. *Biomed Res.*;28(5):261-6.
- <http://www.apitherapy.org> (2001).
- <http://www.bee-hexagon.net> 15 January 2012.
- [http://www.goldinnature.com/apithera pylinks htm](http://www.goldinnature.com/apithera_pylinks.htm). Inter net. 2004.
- Ito S.; Nitta Y.; Fukumitsu H.; Soumiya H.; Ikeno K.; Nakamura T. and Furukawa S. (2012)** Antidepressant-Like Activity of 10-Hydroxy-Trans-2-Decenoic Acid, a Unique Unsaturated Fatty Acid of Royal Jelly, in Stress-Inducible Depression-Like Mouse Model. *Evidence-based complementary and alternative medicine*: 1-6.
- Izuta H.; Chikaraishi Y.; Shimazawa M.; Mishima S. and Hara H. (2009)** 10-Hydroxy-2-decenoic Acid, a Major Fatty Acid from Royal Jelly, Inhibits VEGF-induced Angiogenesis in Human Umbilical Vein Endothelial Cells. *Evidence-based complementary and alternative medicine* 6 (4): 489-494.
- Khattab M.M. (1997)** Bee venom collection as new product from apiaries in Egypt. International symposium of apitherapy, March 8-9th, 1997 of apitherapy center Eldoki. Cairo
- Kim J.; Kim Y.; Yun H.; Park H.; Kim S.Y.; Lee K.G.; Han S. M. and Cho Y. (2010)** Royal jelly enhances migration of human dermal fibroblasts and alters the levels of cholesterol and sphinganine in an in vitro wound healing model. *Nutrition Research and Practice* 4 (5): 362-368.
- Kodai T.; Umebayashi K.; Nakatani T.; Ishiyama K. and Noda N. (2007)** Compositions of royal jelly II. Organic acid glycosides and sterols of the royal jelly of honeybees (*Apis mellifera*). *Chemical & Pharmaceutical Bulletin* 55 (10): 1528-1531.
- Krell R. (1996)** *Value-added products from beekeeping*. FAO Food and Agriculture Organization of the United Nations Roma; 409 pp
- Lercker G.; Caboni M. F.; Vecchi M. A; Sabatini A.G. and Nanetti A. (1993)** Caratterizzazione dei principali costituenti della gelatina reale. *Apicoltura* 8: 27-37.
- Lercker, G. (2003)**. La gelatina reale: composizione, autenticità ed adulterazione. In Atti del Convegno "Strategie per la valorizzazione dei prodotti dell'alveare". Università degli Studi del Molise; Campobasso, 67-81.
- Leung R.; Ho A. and Chan J. (1997)** Royal jelly consumption and hypersensitivity in the community. *Clin Exp Allergy.*; 27:333-336.
- Matsui T.; Yuki Yoshi A.; Doi S.; Sugimoto H.; Yamada H. and Matsumoto K. (2002)** Gastrointestinal enzyme production of bioactive peptides from royal jelly protein and their antihypertensive ability in SHR. *Journal of nutritional biochemistry* 13 (2): 80-86.
- Messia, M.C., Caboni, M.F., Marconi, E. (2005)**. Storage stability assessment of freeze-dried royal jelly by furfural determination. *Journal of agricultural and food chemistry*, 53 (11):4440-4443.
- Mishima S.; Suzuki K.; Isohama Y.; Kuratsu N.; Araki Y.; Inoue M. and Miyata T. (2005)** Royal jelly has estrogenic effects in vitro and in vivo. *J Ethnopharm.*; 101:215-220.
- Münstedt K.; Bargello M.; and Hauenschild A. (2009)** Royal Jelly Reduces the Serum Glucose Levels in Healthy Subjects. *Journal Med Food* 12: 1170-1172.
- Narita Y.; Nomura J.; Ohta S.; Inoh Y.; Suzuki K.M.; Araki Y.; Okada S.; Matsumoto I.; Isohama Y.; Abe K.; Miyata T. and Mishima S. (2006)** Royal jelly stimulates bone formation: physiologic and nutrigenomic studies with mice and cell lines. *Biosci Biotechnol Biochem*;70(10):2508- 2514.
- Norman A.R.; Cicero B.; Eloi S.G.M.; Tariq M.B. and Patricia A. (2011)** "Insect natural products and processes: New treatments for human disease". *Insect Biochemistry and Molecular Biology* 41(10): 747-769,
- Okuda H.; Kameda. K; Morimoto C.; Matsuura Y.; Chikaki M. and Ming J. (1998)** Studies on insulinlike substances in royal jelly and other substances in royal jelly which inhibit angiotensin-converting enzyme. *Honeybee Science* 19 (1): 9-14.
- Petrie A. and Watson P. (1999)** *Statistic for veterinary and animal science*. Blackwell Publishing Company.
- Sabatini, A.G., Marcazzan, G., Caboni, M.F., Bogdanov, S., Almeida-Muradian, L.B. (2009)**

- Quality. and standardisation of royal jelly. *JAAS*, 1: 1-6.
- Scarselli, R., Donadio, E., Giuffrida, M.G., Fortunato, D., Conti, A., Balestreri, E., Felicioli, R., Pinzauti, M., Sabatini, A.G., Felicioli, A. (2005).** Towards royal jelly proteome. *Proteomics*, 5:769-776.
- Schmidt J.O. and Buchmann S.L. (1999)** "Other products of the hive" (In: The hive and the honeybee. Graham, J.M. ed. Dadant & Sons, Hamilton, Illinois, USA. Fourth Printing 952-960.
- Simúth, J. 2001.** Some properties of the main protein of honeybee (*Apis mellifera*) royal jelly. *Apidologie*, 32: 69-80.
- Strum G.; Kranke B. and Rudolph C. (2002).** Rush Hymenoptera venom immunotherapy; a safe and practical protocol for high-risk patients. *J Allergy Clin Immunol* 110(6) : 928-933.
- Terada Y.; Narukawa M. and Watanabe T. (2011)** Specific Hydroxy Fatty Acids in Royal Jelly Activate TRPA1. *Journal of agricultural and food chemistry* 59 (6): 2627-2635.
- Tumanov A.A. and Osipova N.I. (1996)** "Biological determination of traces of substances". Mat. All-Union Conf., 1963, Gorky, USSR, 238-246, 1966. In: Value-added products from beekeeping, Krell, R. (Ed.). FAO Agriculture Services Pulletin. Rome, Italy, pp: 227-240.
- Urtubey N. (2005)** *Apitoxin: from bee venom to apitoxin for medical use*. Termas de Rio Grande Santiago del Estero, Argentina.
- World Health Organization. WHO (2002)** Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction. 4th ed Cambridge, United Kingdom: Cambridge University Press; 1999 (Cited by; Saleh RA. Agarwal A. Oxidative stress and male infertility: From Research Bench to Clinical Practice. *J Androl*; 23 (6):737-752.

الخواص الفيزيوكيميائية والتأثيرات العلاجية لسم النحل والغذاء الملكي المجفدين
أشرف مهدي شروبه ، حسن حسن الطناحي ، حسن نور الدين ، حمدي علي الهنداوي*، سمير يوسف عبدالسيد يوسف

قسم علوم الاغذية- كلية الزراعة – جامعة بنها
*قسم التخدير والألم- كلية الطب البشرى بالاسماعيلية- جامعة قناة السويس

الملخص العربي

تم في هذه الدراسة دراسة التحليل الكيميائي والفيزيائي لسم النحل وغذاء الملكات قبل وبعد اجراء عملية التجفيد لهما ووجد ان عملية التجفيد حسنت كفاءة الانزيمات والمواد الفعالة نتيجة لانخفاض نسبة الرطوبة الى 5% وتم استخدام سم النحل والغذاء الملكي المجفدين لتحسين الكفاءة العلاجية لبعض الأمراض وهي : تحسين الكفاءة الجنسية وتحسين وظائف الكلية وانخفاض نسبة الجلوكوز في الدم حيث تم استخدام جرعات 500 ملي جرام و 1 جرام من المخلوطة (25 ملي عسل+ 500 ملي جرام غذاء ملكات مجفد + 1ملي جرام سم النحل المجفد) / كيلو جرام من وزن الجسم عن طريق الفم في الذكور البالغة من فئران التجارب يوميا لمدة شهر حيث وجد في هذه الدراسة ان معاملة ذكور الفئران البالغين بجرعة 1 جرام من كل المخلوطة السابق / كيلو جرام من وزن الجسم عن طريق الفم يوميا لمدة شهر وحدها نتج عنها زيادة في اوزان الخصية وجسم البربخ وعدد الحيوانات المنوية وهرمون التستوستيرون ونسبة عدد الحيوانات المنوية الحية ومستوى الجلوتاثيون وترافق ذلك مع انخفاض لمستوى المالوندايالديهيد والنسبة المنوية للحيوانات المنوية المشوهة كما اوضحت الدراسة ان معاملة ذكور الفئران بجرعة مقدارها 500 ملي جرام و 1 جرام من المخلوطة السابق / كيلو جرام من وزن الجسم عن طريق الفم يوميا لمدة شهر سببت زيادة تركيز اليوريا في مصل الدم وانخفاضه في البول اضافة الى نقص اليوريا في البول بشكل ملحوظ مقارنة مع مجموعة الكنترول. ان معاملة ذكور الفئران بالمخلوطة السابق بواقع اجرام/ كجم من وزن الجسم عن طريق الفم يوميا لمدة شهر ادت الى انخفاض مستوى الكرياتينين في الدم والبول. كما لوحظ انخفاض نسبة الجلوكوز في الدم في الفئران المعاملة بالمخلوطة السابق بجرعة مقدارها 500 ملي جرام و 1 جرام/ كجم من وزن الجسم لمدة شهر يوميا. كما اوضحت هذه الدراسة زيادة في اوزان فئران التجارب المعاملة بالمخلوطة السابق بواقع اجرام/ كجم من وزن الجسم مقارنة بمجموعة الكنترول. ولهذا توصي هذه الدراسة باستخدام مخاليط سم النحل المجفد والغذاء الملكي المجفد المذابان في عسل النحل في علاج الكفاءة الجنسية والكلية ومرض السكر.